

(12)

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0043

R & T Code 413f0001---01

TECHNICAL REPORT No. 25

Surface Distortions of Asymmetric Line Profiles

by

P. T. Leung and Thomas F. George

Prepared for Publication

in

Chemical Physics Letters

Departments of Chemistry and Physics State University of New York at Buffalo Buffalo, New York 14260

December 1986

Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.

THE FILE COM

A

REPORT DOCUMENTATION PAGE						
14 REPORT SECURITY CLASSIFICATION Unclassified		16. RESTRICTIVE MARKINGS				
24. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited				
26. DECLASSIFICATION/DOWNGRADING SCHEDULE						
4. PERFORMING ORGANIZATION REPORT NUM	BER(S)	S. MONITORING OR	GANIZATION R	EPORT NUMBER(S)		
UBUFFALO/DC/86/TR-25						
Depts. Chemistry & Physics State University of New York	6b. OFFICE SYMBOL (If applicable)	74. NAME OF MONITORING ORGANIZATION				
6c. ADDRESS (City, State and ZIP Code)		76. ADDRESS (City,		le)		
Fronczak Hall, Amherst Campus		Chemistry Program 800 N. Quincy Street				
Buffalo, New York 14260		Arlington, Virginia 22217				
8a. NAME OF FUNDING/SPONSORING ORGANIZATION	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER			MBER	
Office of Naval Research	L	Contract N00014-86-K-0043				
Bc. ADORESS (City, State and ZIP Code) Chemistry Program		10. SOURCE OF FUNDING NOS.		T		
800 N. Quincy Street		PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT	
Arlington, Virginia 22217						
Surface Distortio	ns of Asymmetric	Line Profile	:S			
12. PERSONAL AUTHOR(S) P. T. Leu	ng and <u>Thomas F.</u>	George				
134 TYPE OF REPORT 13b. TIME C		14. DATE OF REPOR			OUNT 13	
16. SUPPLEMENTARY NOTATION	то	Decembe	1 1300		10	
Prepared for publication in Chemical Physics Letters						
17. COSATI CODES	18 SUBJECT TERMS (C. FANO LINE PROF	Continue on reverse if necessary and identify by block number)				
FIELD GROUP SUB. GR.	ASYMMETRIC	FILES METALLIC GRATING DOUBLE-PEAK STRUCTURE				
	ADMOLECULE	SURFACE PLASMON MODE				
Distortions of asymmetric Fano profiles for a molecular system above a metallic grating surface are investigated within a simple model. A double-peak feature as well as a shift of the profile window are observed due to the surface plasmon mode.						
20. DISTRIBUTION/AVAILABILITY OF ABSTRAC		21 ABCTDACT CEC	IRITY CLASSIF	CATION		
	-	21. ABSTRACT SECURITY CLASSIFICATION				
UNCLASSIFIED/UNLIMITED SAME AS RPT.	MOTIC USERS L	Unclassifi			·	
22a. NAME OF RESPONSIBLE INDIVIDUAL		22b. TELEPHONE Nº (Include Area Co	de)	22c. OFFICE SYME	BOL	
Dr. David L. Nelson		(202) 696-4	410	<u> </u>		

SURFACE DISTORTIONS OF ASYMMETRIC LINE PROFILES

P. T. Leung and Thomas F. George
Departments of Chemistry and Physics & Astronomy
State University of New York at Buffalo
Buffalo, New York 14260

Abstract

Distortions of asymmetric Fano profiles for a molecular system above a metallic grating surface are investigated within a simple model. A double-peak feature as well as a shift of the profile window are observed due to the surface plasmon mode.



in the Code of the

I. Int. sduction

Recently, there has been great interest in the study of surface effects on photoabsorption and photodescrption line shapes for an adspecies on a surface. 1-10 This includes the treatment of vibration and vibrationrotation spectra of chemisorbed and physisorbed nolecules. In addition, line profiles involving electronic transitions leading to direct photodissociation of a molecule physisorbed on a rough metallic surface have also been studied for the cases of a metallic spheroid and a shallow sinusoidal grating 7,8 as the substrate. Furthermore, the distortions of the spectra of a dipole on a randomly rough metallic surface have been studied in a fully quantum mechanical approach. Aside from adsorption spectra, 1-9 the line shape of the photodesorption yield has also been examined for physisborbed molecules. 10 However, in most of the previous investigations, the line profile for the free molecule (i.e., in the absence of the surface) has almost always been assumed to be symmetric, often of a Lorentzian type, $^{1-10}$ so that asymmetric distortions are brought about by the presence of the surface. 1-9 One very interesting example is the Langreth line shape which describes the distortion of a vibrational Lorentzian towards a Fano profile, due to energy transfer between the molecule and the surface via the electron-hole-pair mechanism. 3,4 Nevertheless, among these surface distortion phenomena there remains one case uninvestigated, which is that when the line profile for the free molecule case is already asymmetric in nature. This would include, for example, processes like autoionization or predissociation in molecular systems. It is the purpose of the present letter to present a model study of the surface distortions of such "intrinsic" asymmetric profiles. Specifically, we shall study the dipole

photoabsorption spectra on top of a metallic grating, which is considered as a prototype of rough surfaces.

II. Theory

Instead of calculating exactly the line shape function in a quantum mechanical manner, we shall simulate the absorption process by a classical mechanical model and then look for relative profile distortions due to surface effects. This is in the same spirit of the previous work by our group and others in which the free Lorentzian profile for direct photodissociation processes is simulated in the context of the "driven damped oscillator model", and then profile distortions are found by introducing the surface effects into the process via terms including the external driving field and the damping constant.

To describe a "free asymmetric profile", we adopt the mechanical model recently proposed by Sorbello 4 to describe the Fano effect in atomic physics. 11 The model consists of the coupling of a normal mode ($\omega_{_{\scriptsize O}}$) to a lossy broad-band system such as a viscous bath, with both the oscillator and the bath being described by one degree of freedom. The line shape function obtained in this model can be reexpressed 12 in the form

$$I_{o}(\omega) = \frac{(q + \epsilon_{o})^{2}}{1 + \epsilon_{o}^{2}} \sigma_{o}(\omega) , \qquad (1)$$

where q is the asymmetric profile index depending on the <u>ratio</u> of the driving force on the oscillator to that on the bath, and $\sigma_{0}(\omega)$ is the background intensity due to direct excitation of the bath. In case of actual molecular processes, q is defined as the <u>ratio</u> of the transition matrix elements from the initial state to two states whose linear combination gives the final state of the system, 11,13 and $\sigma_{0}(\omega)$ corresponds

to the cross section for direct excitation of the continuum. The reduced energy variable ϵ_0 in Eq. (1) is expressed as

$$\varepsilon_{0} = \frac{2(\omega - \omega_{0} - \Delta\omega_{0})}{\gamma_{0}} , \qquad (2)$$

where $\Delta\omega_0$ and γ_0 are the level shift and decay rate of the system, respectively. In general, $\Delta\omega_0$ << ω_0 , and hence

$$\varepsilon_{0} = \frac{2\Delta\omega}{\gamma_{0}}$$
(3)

with $\Delta \omega \equiv \omega - \omega_0$.

To investigate the distortions of $I_0(\omega)$ due to the presence of a rough metallic surface, we shall consider the oscillator as a point dipole oriented perpendicularly on top of a shallow sinusoidal grating, a case which we have formulated before for a Lorentzian $I_0(\omega)$. Analogously, we have the surface line shape function expressed as

$$I(\omega) = \frac{(q + \varepsilon)^2}{1 + \varepsilon^2} |\hat{n}_{\mu} \cdot [1 + A] \cdot \hat{n}|^2 \sigma(\omega) , \qquad (4)$$

where quantities without the subscript 'o' refers to those at the surface, and ϵ is now defined as

$$\varepsilon = \frac{2\Delta\omega}{\gamma} \quad . \tag{5}$$

We have in this approach neglected the surface effects on q and ω_0 ; this is justified since, for the former, the effects enter into the <u>ratio</u> of two transition matrix elements which may be regarded as approximately cancelling each other, and for the latter such effects have been found to be small. ¹⁴
In this way, surface effects will only enter via the decay rate γ in Eq. (5)

and the field enhancement factor A in Eq. (4), the latter including both of the Fresnel reflected and the surface plasmon fields. 7 \hat{n}_{μ} and \hat{n} in Eq. (4) are the unit vectors of the molecular dipole and the incident field, respectively. For a dipole located at (0,0,d), it has been found that 7

$$|\hat{\mathbf{n}}_{u}\cdot[1+A]\cdot\hat{\mathbf{n}}|^{2} = |1+Re^{2ik_{z}d}+Se^{(ik_{z}-\Gamma_{g})d}|^{2}\sin^{2}\theta$$
, (6)

where θ is the angle of incidence of the laser light, and the quantities R, S, $\mathbf{k}_{\mathbf{Z}}$ and $\Gamma_{\mathbf{g}}$ are defined as in Ref. 7. Furthermore, the surface-induced molecular decay rate can be expressed as

$$\frac{Y}{Y_0} = 1 + \frac{3}{2} \frac{q}{k^3} \text{ Im } G^F \left(1 + \frac{\text{Im } G^R}{\text{Im } G^F}\right) ,$$
 (7)

where again all symbols are the same as in Ref. 7.

With the results in Eqs. (6) and (7) substituted into Eqs. (4) and (5), we can calculate the line profile for a molecule on top of a metallic grating by assuming some reasonable form for $\sigma_{0}(\omega)$. Within the present model study, we have considered the following three different cases with the surface always taken to be a sinusoidal silver grating whose dielectric constant as a function of photon energy is available in a numerically-fitted form. ⁷

Case (i) $\sigma_{O}(\omega) = \sigma(\omega) = constant$.

In this case, $I_o(\omega)$ may simulate certain atomic autoionization processes (e.g., the photoabsorption line shape of helium in the neighborhood of the first $^1P^0$ resonance). 11,15 In Fig. 1, we show both the line shapes $I_o(\omega)$ and $I(\omega)$ for some hypothetical profile constants (see the figure caption). We see that the frequency corresponding to the zero of the

absorption cross section shifts towards a larger value. This is because q < 0 in this case and $\gamma > \gamma_0$ as in Eq. (7).

Case (ii)
$$\sigma_o(\omega) = \sigma(\omega) = 1/\omega^2$$
.

In this case, $I_o(\omega)$ simulates a molecular predissociation process (e.g., the predissociation of the $D^1\Pi^+_u$ levels by the $B^{*1}\Sigma^+_u$ continuum in the photodissociation experiment of hydrogen molecules 16). The result is shown in Fig. 2, where we see that the 'window' at the steep edge at the low frequency end of the original profile is broadened due to the enhancement effect by the surface plasmon field. At the high frequency end of the distorted profile, the surface plasmon resonance leads to a new window.

Case (iii)
$$\sigma_o(\omega) = \frac{\gamma_o}{(\omega - \omega_o)^2 + \gamma_o^2/4}$$

For the above Lorentzian, $\sigma(\omega) \neq \sigma_O(\omega)$ since γ_O is replaced by γ in Eq. (7). The effects of the surface distortion of the profile $I_O(\omega)$ are similar to case (ii) (Fig. 3), except that the original window is not broadened as much.

III. Results and Discussion

In all cases (i) to (iii), the dipole is fixed at a distance of 500 Å from the grating, which has a spatial period of 8000 Å and an amplitude-to-period ratio of approximately 10⁻². In addition, the angle of incidence is fixed at 0.6 rad and the surface plasmon resonance occurs at about 2.9 eV photon energy. We observe that the "double peak" feature appears in all cases, corresponding to the resonance of the molecular system and that of the surface plasmon, respectively. Furthermore, a surface enhancement effect is in general observed, implying that enhanced molecular photopredissociation may also be possible, provided the molecule is not located

too close to the surface and the resonant plasmon field decays very slowly in the direction perpendicular to the surface. 7,17

Acknowledgments


One of us (P. T. L.) wishes to acknowledge Professor Y. C. Lee and Dr. Daniel A. Jelski for very useful discussions. This research was supported by the Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract F49620-86-C-0009, the Office of Naval Research, and the National Science Foundation under Grant CHE-8519053. The United States Government is authorized to reproduce and distribute reprints notwithstanding any copyright notation hereon.

References

- 1. H. Metiu and W. E. Palke, J. Chem. Phys. 69, 2574 (1978).
- 2. J. W. Gadzuk and A. C. Luntz, Surf. Sci. 144, 429 (1984).
- 3. D. C. Langreth, Phys. Rev. Lett. <u>54</u>, 126 (1985).
- 4. R. S. Sorbello, Phys. Rev. B <u>32</u>, 6294 (1985).
- 5. J. E. Adams, J. Chem. Phys. <u>84</u>, 3589 (1986).
- A. Nitzan and L. E. Brus, J. Chem. Phys. <u>74</u>, 5321 (1981); P. Das and H. Metiu, J. Phys. Chem. <u>89</u>, 4680 (1985).
- 7. P. T. Leung and T. F. George, J. Chem. Phys. <u>85</u>, 4729 (1986).
- 8. E. Hood, P. K. Aravind and H. Metiu, Surf. Sci. <u>109</u>, 95 (1981).
- 9. D. Agassi, Phys. Rev. B <u>33</u>, 3873 (1986).
- 10. Z. W. Gortel, P. Piercy, R. Teshima and H. J. Kreuzer, Surf. Sci. 165, L12 (1986).
- 11. U. Fano, Phys. Rev. 124, 1866 (1961).
- 12. We have changed some of Sorbello's original notations from Ref. 4. Here we denote every quantitiy which refers to the free molecule case in the absence of the surface with the subscript 'o'.
- 13. T. F. O'Malley and S. Geltman, Phys. Rev. <u>137</u>, A1344 (1965).
- 14. See, e.g., R. R. Chance, A. Prock and R. Silbey, Adv. Chem. Phys. <u>37</u>, 1 (1978).
- P. G. Burke and D. D. McVicar, Proc. Phys. Soc. 86, 989 (1965).
- M. Glass-Maujean, P. M. Guyon and J. Breton, Phys. Rev. A <u>33</u>, 346 (1986).
- 17. T. J. Chuang, Surf. Sci. Rep. 3, 1 (1982).

Figure Captions

- 1. Distortion of the Fano profile with $\sigma_{0}(\omega)$ = constant. The profile constants are: $\gamma_{0} = 5 \times 10^{2}$, $\omega_{0} = 1.6 \times 10^{4}$ and q = -2.65. Note that the scales for $I_{0}(\omega)$ and $I(\omega)$ are different. The y-axis quantities are in arbitrary units for all three figures.
- 2. Distortion of the Fano profile with $\sigma_{o}(\omega) = 1/\omega^{2}$. The profile constants are: $\gamma_{o} = 4 \times 10^{3}$, $\omega_{o} = 1.17 \times 10^{4}$ and q = +3.0.
- 3. Distortion of the Fano profile with a Lorentzian $\sigma_0(\omega)$. The profile constants are the same as in Fig. 2.






Fig. 2




Fig. 3

01/1113/86/2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	No. Coples	•	No. Copies
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401 Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

Dr. J. E. Jensen Hughes Research Laboratory 3011 Malibu Canyon Road Malibu, California 90265

Dr. J. H. Weaver
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. A. Reisman Microelectronics Center of North Carolina Research Triangle Park, North Carolina 27709

Dr. M. Grunze Laboratory for Surface Science and Technology University of Maine Orono, Maine 04469

Dr. J. Butler Naval Research Laboratory Code 6115 Washington D.C. 20375-5000

Dr. L. Interante Chemistry Department Rensselaer Polytechnic Institute Troy, New York 12181

Dr. Irvin Heard Chemistry and Physics Department Lincoln University Lincoln University, Pennsylvania 19352

Or. K.J. Klaubunde Department of Chemistry Kansas State University Manhattan, Kansas 66506 Dr. C. B. Harris Department of Chemistry University of California Berkeley, California 94720

Dr. F. Kutzler
Department of Chemistry
Box 5055
Tennessee Technological University
Cookesville, Tennessee 38501

Dr. D. Dilella Chemistry Department George Washington University Washington D.C. 20052

Dr. R. Reeves Chemistry Department Renssaeler Polytechnic Institute Troy, New York 12181

Dr. Steven M. George Stanford University Department of Chemistry Stanford, CA 94305

Dr. Mark Johnson Yale University Department of Chemistry New Haven, CT 06511-8118

Dr. W. Knauer Hughes Research Laboratory 3011 Malibu Canyon Road Malibu, California 90265

Or. G. A. Somorjai Department of Chemistry University of California Berkeley, California 94720

Dr. J. Murday
Naval Research Laboratory
Code 6170
Washington, D.C. 20375-5000

Dr. J. B. Hudson Materials Division Rensselaer Polytechnic Institute Troy, New York 12181

Dr. Theodore E. Madey
Surface Chemistry Section
Department of Commerce
National Bureau of Standards
Washington, D.C. 20234

Dr. J. E. Demuth
IBM Corporation
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. M. G. Lagally
Department of Metallurgical
and Mining Engineering
University of Wisconsin
Madison, Wisconsin 53706

Dr. R. P. Van Duyne Chemistry Department Northwestern University Evanston, Illinois 60637

Dr. J. M. White Department of Chemistry University of Texas Austin, Texas 78712

Dr. D. E. Harrison Department of Physics Naval Postgraduate School Monterey, California 93940 Dr. R. L. Park
Director, Center of Materials
Research
University of Maryland
College Park, Maryland 20742

Dr. W. T. Peria Electrical Engineering Department University of Minnesota Minneapolis, Minnesota 55455

Dr. Keith H. Johnson
Department of Metallurgy and
Materials Science
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. S. Sibener
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Arnold Green Quantum Surface Dynamics Branch Code 3817 Naval Weapons Center China Lake, California 93555

Dr. A. Wold Department of Chemistry Brown University Providence, Rhode Island 02912

Dr. S. L. Bernasek Department of Chemistry Princeton University Princeton, New Jersey 08544

Dr. W. Kohn Department of Physics University of California, San Diego La Jolla, California 92037

Dr. F. Carter Code 6170 Naval Research Laboratory Washington, D.C. 20375-5000

Or. Richard Colton Code 6170 Naval Research Laboratory Washington, D.C. 20375-5000

Dr. Dan Pierce National Bureau of Standards Optical Physics Division Washington, D.C. 20234

Dr. R. Stanley Williams
Department of Chemistry
University of California
Los Angeles, California 90024

Dr. R. P. Messmer Materials Characterization Lab. General Electric Company Schenectady, New York 22217

Dr. Robert Gomer
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Or. Ronald Lee R301 Naval Surface Weapons Center White Oak Silver Spring, Maryland 20910

Dr. Paul Schoen Code 6190 Naval Research Laboratory Washington, D.C. 20375-5000 Dr. John T. Yates Department of Chemistry University of Pittsburgh Pittsburgh, Pennsylvania 15260

Dr. Richard Greene Code 5230 Naval Research Laboratory Washington, D.C. 20375-5000

Dr. L. Kesmodel
Department of Physics
Indiana University
Bloomington, Indiana 47403

Dr. K. C. Janda University of Pittsburg Chemistry Building Pittsburg, PA 15260

Or. E. A. Irene
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. Adam Heller Bell Laboratories Murray Hill, New Jersey 07974

Dr. Martin Fleischmann Department of Chemistry University of Southampton Southampton 509 5MH UNITED KINGDOM

Dr. H. Tachikawa Chemistry Department Jackson State University Jackson, Mississippi 39217

Dr. John W. Wilkins Cornell University Laboratory of Atomic and Solid State Physics Ithaca, New York 14853

Dr. R. G. Wallis Department of Physics University of California Irvine, California 92664

Dr. D. Ramaker Chemistry Department George Washington University Washington, D.C. 20052

Dr. J. C. Hemminger . Chemistry Department University of California Irvine, California 92717

Dr. T. F. George Chemistry Department University of Rochester Rochester, New York 14627

Or. G. Rubloff IBM Thomas J. Watson Research Center P.O. Box 218 Yorktown Heights, New York 10598

Dr. Horia Metiu Chemistry Department University of California Santa Barbara, California 93106

Dr. W. Goddard
Department of Chemistry and Chemical
Engineering
California Institute of Technology
Pasadena, California 91125

Dr. P. Hansma Department of Physics University of California Santa Barbara, California 93106

Dr. J. Baldeschwieler
Department of Chemistry and
Chemical Engineering
California Institute of Technology
Pasadena, California 91125

Dr. J. T. Keiser Department of Chemistry University of Richmond Richmond, Virginia 23173

Dr. R. W. Plummer
Department of Physics
University of Pennsylvania
Philadelphia, Pennsylvania 19104

Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland, Ohio 41106

Dr. N. Winograd
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Dr. Roald Hoffmann Department of Chemistry Cornell University Ithaca, New York 14853

Or. A. Stecki
Department of Electrical and
Systems Engineering
Rensselaer Polytechnic Institute
Troy, NewYork 12181

Dr. G.H. Morrison Department of Chemistry Cornell University Ithaca, New York 14853

EMD D

2 - 8 7

DTIC